

MONTE CARLO CALCULATIONS OF TWO-DIMENSIONAL COMBUSTOR FLOWS

Final Report

Stephen B. Pope

4/12/82

U.S. Army Research Office
DAA929 81 K 0031

Massachusetts Institute of Technology

Cambridge, MA 02139



Approved for public release; distribution unlimited.

81 97 . 1 037

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
T REPORT NUMBER		3 RECIPIENT'S CATALOG NUMBER
	AD-A113 78	
4 TITLE (and Subtitie)		5 TYPE OF REPORT & PERIOD COVERED Final Report
MONTE CARLO CALCULATIONS OF TWO-DIMENSIONAL		1/13/81 - 1/12/82
COMBUSTOR FLOWS		6 PERFORMING ORG REPORT NUMBER
7. AUTHOR(e)		B CONTRACT OR GRANT NUMBER(*)
Stephen B. Pope		DAA929-81-K-0031
Stephen B. Tope		
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10 PROGRAM ELEMENT, PROJECT, YASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
U. S. Army Research Office		4/12/82
Post Office Box 12211 Research Triangle Park, NC 27709		13 NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS(II dille	rent from Controlling Office)	15 SECURITY CLASS. (of this report)
	•	
Massachusetts Institute of Technology Cambridge, MA 02139		United Street
Campfinge, Fa 02133		154. DECLASSIFICATION DOWNGRADING SCHEDULE
Approved for public release; o		
17. DISTRIBUTION STATEMENT (of the abatract enter	red in Block 20, it different in	же керип)
18. SUPPLEMENTARY NOTES The view, opinions, and/or fin author(s) and should not be coposition, policy, or decision, 19. KEY WORDS (Continue on reverse side if necessar	nstrued as an off unless so designa	ated by other documentation.
Combustion, Turbulence, Monte C		
40. ABSTRACY (Continue on reverse elds If necessary	and identify by block number	
Progress has been made in develo properties of two-dimensional tu	ping a computation rbulent reactive feld is calculated thod is used to so fractions. In teree Monte Carlong has been develo	al procedure to calculate the low fields appropriate to gas by a standard finite-difference lve the transport equation for he course of this project, a method has been devised and an ped. For plane constant-densit

DD 17000 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

FOREWORD

"Monte Carlo Calculations of Two-Dimensional Combustor Flows" was originally a two-year research program being performed at the Massachusetts Institute of Technology. Since the principal investigator left MIT at the end of 1981, the contract was terminated, and the research is being continued at Cornell University under contract number DAA929-82-K-0017. This final report is for the one-year period 1/13/81 - 1/12/82.



Accession For	7
NTIS GRANT DTIC TAB Unamounced Justification	
Ву	
Distribution/ Availability Codes	
Avnil and/or Dist Special	
H	į

Problem Statement

The objective of this work was to develop and demonstrate a combined Monte Carlo/finite-difference procedure for calculating the properties of two-dimensional combustor flows.

It is a formidable theoretical problem to calculate the turbulent reacting flow properties in a gas turbine combustor. Added to the well-known problems of calculating any turbulent flow, are the difficulties caused by the highly non-linear combustion reactions. An approach that goes a long way to overcoming the problems caused by the non-linear reactions is to describe the turbulent reacting flow field in terms of the joint probability density functions (pdf) of the species mass fractions. The transport equation for the joint pdf of the species contains an exact term pertaining to reaction. That is, irrespective of the complexity and non-linearity of the reaction scheme, in the joint pdf equation the reaction term does not have to be modelled. For this reason, several authors have used this equation to calculate turbulent reactive flows: Dopazo (1975), Janicka, Kolbe and Kollmann (1979) and Pope (1981a).

By standard techniques (e.g. finite-differences), the joint pdf equation would be extremely expensive to solve numerically (if, ineeed, it were possible), except in simple cases. However, by using a Monte Carlo method, the equation can be solved at a reasonable cost for situations of practical complexity. The basic Monte Carlo method is described by Pope (1981b).

The Monte Carlo solution of the joint pdf equation deals only with the species fields. The mean velocity and turbulence fields are calculated by a standard finite difference procedure using the $k-\epsilon$ turbulence model.

Results

In the Monte Carlo method described by Pope (1981b), the joint pdf is represented by ensembles of elements at discrete grid nodes. There is both a truncation error and a statistical error associated with the discretization of physical space. An improved method has been developed in which the elements are distributed throughout the flow field — not just at discrete nodes. This method resembles a Lagrangian particle tracking procedure with a random walk to simulate turbulent diffusion. The numerical error is greatly reduced, and hence the computational economy is improved. The method is described by Pope (1982a).

This method has been implemented for plane, constant-density flows. The two major tasks were to treat the boundaries correctly and to develop the necessary least-squares cubic spline procedures. The cubic splines are used to extract information about mean quantities from the elements randomly distributed through the flow field. At the end of the contract, the code had been developed and testing was under way.

One of the components of the Monte Carlo method is a model of turbulent mixing. Most investigators use Curl's (1963) model or a variant of it (Dopazo, 1979; Janicka, Kolbe and Kollmann, 1979). In the course of this investigation it was found that these models have serious defects. Specifically, when the models are applied to the simple case of decaying fluctuations they should predict a Gaussian pdf. However, the predicted pdf is far from Gaussian and, in fact, its fourth and higher even moments are infinite.

Because of this observed defect, a new class of mixing models has been developed that overcomes this problem (Pope, 1982b). The improved models result in approximately Guassian pdf's, and a consequent gain in accuracy is expected in the pdf calculations.

Personnel and Publications

The principal investigator Stephen B. Pope and the graduate student Babak Minaie have worked on the project throughout 1981. The following papers have been prepared for publication:

"An Improved Turbulent Mixing Model," by S. B. Pope, to be published in Comb. Sci. Technol., 1982.

"PDF Methods in Turbulent Reactive Flows," by S. B. Pope (in preparation), to be published in Prog. Energy Comb. Sci., 1982.

Bibliography

Curl, R. L. (1963) A. I. Ch. E. J., 9, 175.

Dopazo, C. (1975) Phys. Fluids, 18, 397.

Dopazo, C. (1979) Phys. Fluids, 22, 20.

Janicka, J., Kolbe, W. and Kollmann, W. (1979) J. Nonequilib. Thermodyn., 4, 47.

Pope, S. B. (1981a) <u>Eighteenth Symp. (Int'1.) on Combustion</u>, 1001, Combustion Institute.

Pope, S. B. (1981b) <u>Combust. Sci. Technol.</u>, <u>25</u>, 159.

Pope, S. B. (1982a) Prog. Energy Comb. Sci. (in preparation).

Pope, S. B. (1982b) Combust. Sci. Technol., to be published.